

## A CAVE OBLIGATE AMPHIPOD

---


### TAXONOMY

**Scientific name:** *Stygobromus quatsinensis* (Holsinger and Shaw, 1987)

**Common name:** Unnamed; a cave obligate amphipod scud

**Taxonomic comments:**

Recently described species (1987) on Vancouver Island, British Columbia. All members of the genus are stygobiont (obligate to subterranean waters). Closely related to other members of the phylogenetic hubbsi group but discovery of this species may cause revision of the physical characters used to define the group (see Holsinger and Shaw 1987).



### DESCRIPTION

**Basic description:** A cave obligate amphipod.

**General description:**

Blind, unpigmented subterranean amphipod. Medium sized groundwater species distinguished by presence of 2 or 3 submarginal setae on distal part of posterior margin of propod of gnathopod 1, inserted below defining angle; and absence of distal peduncular process on uropod 1 of male (Wang and Holsinger, 2001). Largest female 7 mm, largest male 6.3 mm; male and female generally similar except for slight differences in gnathopod and uropod appendages (Holsinger and Shaw 1987).

**Ecology:**

Occurrence on Coronation Island, southeastern Alaska, represents the western hemisphere high latitude record for any cave adapted species (Carlson 1996). Method of origination in current locations is unknown; apparently occurs only on coastal islands such as Vancouver Island, BC, which has been isolated from mainland North America for around 40 million years. Known distribution may be the result of subterranean communities remaining stable through periods of glacial coverage (cave-adapted fauna are perhaps the only invertebrates that could survive glaciation in situ) or radiation from existing southern mainland populations after glaciers receded (Holsinger and Shaw 1987, Carlson 1997a).

**Food:**

Thought to be a predator-scavenger of organic matter that falls below the surface of cave drip pools and streams; uses mouth appendages (gnathopods) to capture food (Carlson 1997b).

**Habitat:**

Occurs in resurgence areas of limestone caves or karst formations; in cave streams, springs or drip pools with substrates of mud, pebble, cobble or bare rock. Water bodies characterized by small amounts of organic matter, low temperatures (3.0 to 8.5°C) and pH of approximately 7.5-8.0 (Holsinger and Shaw 1987, Carlson 1997a, Carlson 1997c). Organic matter enters karst system in

runoff, seeps and streams, is broken down by microscopic organisms, and provides nutrients and energy to cave inhabitants (Carlson 1997b).

## STATUS

**Global rank:** G2G3 (1999-12-02)

**Global rank reasons:**

Global rank reasons currently unavailable.

**State rank:** S2S3 (2004-08-04)

**State rank reasons:**

A recently described species; suspected rare. Known only from karst formations on coastal islands adjacent to Prince of Wales Island in the Alexander Archipelago; distribution information is likely incomplete. Sensitive to disturbance of karst habitat by human activity which could alter water infiltration rates, sediment production and debris transport, and water chemistry.

## DISTRIBUTION AND ABUNDANCE

**Range:**

**Global range:**

Caves on Vancouver Island, British Columbia, and elsewhere in karst groundwater habitats in the Alexander Archipelago of southeastern Alaska (Wang and Holsinger, 2001).

**State range:**

In Southeast Alaska, occurs in karst groundwater habitats in the Alexander Archipelago, including Nautilus cave, Heceta Island, and in numerous springs and caves on Dall, Baker, Sumez and Coronation Islands adjacent to Prince of Wales Island (Carlson 1994, 1996, 1997a, Holsinger et al. 1997).

**Abundance:**

**Global abundance:**

Unknown. See State abundance comments below.

**State abundance:**

Unknown. A study of aquatic and terrestrial cave-associated invertebrates in Southeast Alaska found overall species abundance and richness decreased as collection sites progressed from east to west and from north to south (i.e., higher species abundance closer to Prince of Wales Island and mainland Alaska); overall abundance per collection site was low (<10 organisms) (Carlson 1997a).

**Trends:**

**Global trend:**

Unknown.

**State trend:**

Unknown.

## EXISTING PROTECTION

### Global protection:

See State protection below.

### State protection:

In the US, the Federal Cave Resources Protection Act of 1988 safeguards caves on federal lands by regulating use, requiring permits for removal of cave resources, prohibiting destruction, and allowing for cave locations to remain confidential and requiring that caves be included in land management plans (New Mexico Center for Wildlife Law 1998). Directed by this act, the Tongass National Forest in Southeast Alaska included standards and guidelines for cave and karst management in the 1997 Tongass Forest Plan Revision Final EIS, which includes assessment of “karst vulnerability” for karst resources to any land use (USFS 2002). Around 467,600 acres of soluble rock (where karst formations occur, usually in limestone and marble) underlie lands currently administered by the Tongass National Forest. Fifty-three percent of that acreage is managed under a Wilderness Group or Natural Setting Land Use Designation (LUD) and is protected from development or resource harvest. The remaining 47 percent is in Development LUDs, and more than half of this area is mapped as suitable for timber harvest. However, through “karst vulnerability” assessment, more acreage may eventually be managed protectively (USFS 2002).

## CHALLENGES

### Global challenges:

See State challenges below.

### State challenges:

Cave-adapted species are perhaps the most sensitive of all invertebrates to disturbance or impact; their extremely specific habitat requirements and physiological traits make them unable to compete with terrestrial/surface-dwelling invertebrates (Carlson 1997b). One of the largest challenges regarding this species is disturbance of cave or karst habitat, especially by human activity. Timber harvest, road building and other development can affect water infiltration rates, sediment production and debris transport, and introduce pollutants or organic materials which can alter water chemistry (USFS 2002). Temperature changes in aquatic environments resulting indirectly from anthropogenic effects are another concern and may be the greatest threat to aquatic organisms geographically constrained to certain karst areas (Carlson, pers. comm.). Anthropogenically introduced non-native invertebrate species (including the collembolan *Willowsia* and Formicid ants) can also threaten cave-dwelling communities (Carlson 1997a).

## RESEARCH AND INVENTORY NEEDS

### Global research needs:

See State research needs below.

### State research needs:

Research to determine life history and specific habitat requirements should be a priority. Genetic studies of local populations on Vancouver Island and Southeast Alaska should be conducted to help

establish dispersal timing and provide important information on colonization of offshore islands (Shaw and Davis 2000).

**Global inventory needs:**

See State inventory needs below.

**State inventory needs:**

Recent description of this species and its distribution suggests new occurrences of *S. quatsinensis* may be found in the future; further exploration of freshwater cave and karst aquatic habitats should be undertaken to establish full geographic range. Catalogue and inventory all known occurrences.

## **CONSERVATION AND MANAGEMENT NEEDS**

**Global conservation and management needs:**

See State conservation and management needs below.

**State conservation and management needs:**

Conservation of cave and karst habitats should be a high priority. Careful environmental and faunal studies before and after timber harvest could help assess degree of threat as a result of habitat alteration.

## LITERATURE CITED

- Carlson, K. R. 1994. Inventory and assessment of ecological relationships between cavernicolous (cave-associated) invertebrate species and their interactions in representative karst ecosystems on carbonate terrain in the Ketchikan area Tongass National Forest, Part I: Dall Island. Karst Biosciences. Middletown, MD.
- Carlson, K. R. 1996. Inventory and assessment of ecological relationships between cavernicolous (cave-associated) invertebrate species and their interactions in representative karst ecosystems on carbonate terrain in the Ketchikan area Tongass National Forest, Part II: Coronation Island. Karst Biosciences. Middletown, MD.
- Carlson, K. R. 1997a. The distribution of troglobitic and troglophilic invertebrates in southeast Alaska. Proceedings of the 1997 Karst and Cave Management Symposium/13th National Cave Management Symposium, October 7-10. Bellingham, WA. Pp. 28-33.
- Carlson, K. R. 1997b. Text for Ketchikan Radio Broadcast 1997: a brief introduction to invertebrate life in southeast Alaskan caves. Unpublished text. Karst Biosciences. Middletown, MD.
- Carlson, K. R. 1997c. Invertebrate habitat complexity in southeast Alaskan karst ecosystems. Proceedings of the 1997 Karst and Cave Management Symposium/13th National Cave Management Symposium, October 7-10. Bellingham, WA. Pp. 34-43.
- Carlson, K. R. 2004. Personal communication of review comments regarding threats to *Stygobromus quatsinensis*, to Jodi McClory, Research Assistant, Alaska Natural Heritage Program, University of Alaska, Anchorage, AK.
- Holsinger, J. R. and D. P. Shaw. 1987. *Stygobromus quatsinensis*, a new amphipod crustacean (Crangonyctidae) from caves on Vancouver Island, British Columbia, with remarks on zoogeographic relationships. Canadian Journal of Zoology 65:2202-2209.
- Holsinger, J. R., K. R. Carlson and D. P. Shaw. 1997. Biogeographic significance of recently discovered amphipod crustaceans (*Stygobromus*) in caves of southeastern Alaska and Vancouver Island. Proceedings of the 12<sup>th</sup> International Congress of Speleology, 1997, Switzerland. Symposium 9:347-349.
- New Mexico Center for Wildlife Law. 1998. Federal cave resources protection act: summary from the Federal Wildlife Laws Handbook. University of New Mexico (UNM) Institute of Public Law (IPL). <http://ipl.unm.edu/cwl/fedbook/fedcave.html>. Accessed 7/30/04.
- Shaw, D. P. and M. Davis. 2000. Invertebrates from caves on Vancouver Island. Pp. 121-124 In: Darling, L. (ed) Proceedings of a conference on the biology and management of species and habitats at risk, Kamloops, BC 15-19 Feb. 1999. B.C. Ministry of Environment, Lands, and Parks, Victoria, B.C. and University of the Cariboo, Kamloops, B.C. 490 pp.

U.S. Forest Service (USFS). 2002. Physical and biological environment, karst, affected environment, karst 3-X draft seis. Chapter 3 In: Draft supplemental environmental impact statement, roadless area evaluations for wilderness recommendations. Tongass Land Management Plan Revision. U.S. Forest Service Vol 1:Draft SEIS. Juneau, AK.

Wang, D. and J.R. Holsinger. 2001. Systematics of the subterranean amphipod genus *Stygobromus* (Crangonyctidae) in western North America, with emphasis on species of the hubbsi group. *Amphipacifica*, 3(2): 39-147.

---

### Acknowledgements

**State Conservation Status, Element Ecology & Life History Author(s):** Gotthardt, T.A. and J.G. McClory  
**State Conservation Status, Element Ecology & Life History Edition Date:** 23Mar2005

**Reviewer(s):** Kent R. Carlson, Karst Biosciences, Middletown, MD.

Life history and Global level information were obtained from the on-line database, NatureServe Explorer ([www.natureserve.org/explorer](http://www.natureserve.org/explorer)). In many cases, life history and Global information were updated for this species account by Alaska Natural Heritage Program zoologist, Tracey Gotthardt. All Global level modifications will be sent to NatureServe to update the on-line version.

**Global NatureServe Conservation Status Factors Edition Date:** 24Jun2004

**Global NatureServe Conservation Status Factors Author:** Cordeiro, J.

**Global Element Ecology & Life History Edition Date:** 24Jun2004

**Global Element Ecology & Life History Author(s):** Cordeiro, J.

---